

# Potential for novel food products from agroforestry trees: A review

R.R.B. Leakey (1999)

*Institute of Terrestrial Ecology, Bush Estate, Penicuik, Midlothian, Scotland, UK, EH26 0QB  
R.Leakey@ite.ac.uk*

The domestication of trees for agroforestry approaches to poverty alleviation and environmental rehabilitation in the tropics depends on the expansion of the market demand for their non-timber forest products. This paper reviews published data on the nutritive values of the flesh, kernels and seedoils of the seventeen fruit tree species that have been identified, in four ecoregions of the tropics, by subsistence farmers as their top priorities for domestication.

In some species, genetic variation in nutritive value has been reported, but in most species there is still inadequate information on which to base programmes for the genetic improvement of these species. Farmers and agroforesters have identified many of the biological constraints relevant to their viewpoint on production, but there is a need for inputs from the food industry into the identification of the desirable traits and characteristics of potentially novel food products. This paper calls for greater collaboration between agroforesters and the food industry in the effort to promote the domestication and commercialization of underutilized tree products.

**Keywords:-** Domestication, West Africa, Humid forest, Sahel, Miombo Woodlands, Amazonia, *Irvingia gabonensis*, *Dacryodes edulis*, *Ricinodendron heudelottii*, *Chrysophyllum albidum*, *Garcinia kola*, *Adansonia digitata*, *Vitellaria paradoxa*, *Parkia biglobosa*, *Tamarindus indica*, *Zizyphus mauritiana*, *Sclerocarya birrea*, *Uapaca kirkiana*, *Zizyphus mauritiana*, *Vangueria infausta*, *Azanza garckeana*, *Inga edulis* and *Bactris gasipaes*.

## INTRODUCTION

New initiatives in agroforestry are seeking to promote poverty alleviation and environmental rehabilitation in Developing Countries, through the integration of indigenous trees, whose products have traditionally been gathered from natural forests, into tropical farming systems (ICRAF, 1997). This is being done in order to provide marketable products from farms that will generate cash for resource-poor rural and peri-urban households. One important component of this approach is the domestication of the local tree species that have commercial potential in local, regional or even international markets (Leakey & Simons, 1997). Consequently, in collaboration with ICRAF (International Centre for Research in Agroforestry), farmers in four ecoregions of the tropics (the humid and dry zones of West Africa, Amazonia, and southern Africa) have identified their priority indigenous trees for “domestication”, from among the many that have been, and are still being, used traditionally, to provide people’s needs for food and nutritional security (Plate 1). For most of these hitherto wild species, little attention has been made to seek market opportunities for the products within the international food industry, although in some instances research has been carried out to assess their food

value and potential for domestication. To meet the objective of market creation and expansion, it is important to identify potential market niches and then to determine whether there are important product characteristics, which should be improved through genetic selection. While some traits that are relatively easy to identify do benefit the farmer, there are undoubtedly others that are important to the food industry, but that require more sophisticated evaluation in collaboration with the private sector. This review seeks to draw together, in priority order for each region, all the existing information on the characteristics of the products from the tree species that farmers have identified as being their preferred choice for domestication.

## HUMID LOWLANDS OF WEST AFRICA

### 1. *Irvingia gabonensis* (O'Rorke) Baill. and related species (Bush Mango or Dika Nut).

This fruit is like that of a small, cultivated Mango in appearance (Plate 2); although they are unrelated. The pulp of this fruit is eaten fresh and the kernel of the nut is a food additive. The flesh is juicy and varies between sweet and bitter. The sweeter form is generally considered to be *I. gabonensis* var. *gabonensis*, while the bitter form is var. *excelsa*, now called *Irvingia wombolu*. Trees are being selected for the sweetness of their fruits, fruit size, colour and other desirable traits (Ladipo *et al.*, 1996), but not so far for any kernel traits. The pulp can be used for the preparation of juice, jelly and jam. The extraction rate of juice from the fruit pulp was 75% and the sugar concentration of this juice is comparable with pineapples and oranges (Akubor, 1996), but with a higher ascorbic acid content (67mg/100 ml). This concentration of ascorbic acid is also nearly three times that of *Dacryodes edulis* and *Chrysophyllum albidum* (Achinewhu, 1983). These fruits are therefore a good local source of vitamin A.

Evaluation of the wine making potential of the juice (Akubor, 1996), found that wine produced after 28 days fermentation had 8.12% alcohol content. Sensory evaluation showed no significant difference in colour, mouthfeel, sweetness, flavour and general acceptability from a German reference wine.

A study of fruit ripening and storage (Joseph & Aworh, 1991) has shown that fruits harvested at the mature green stage and ripened at 26-29°C were preferred to tree-ripened fruits in colour and texture, although they were both comparable in composition. Fruits held at 12-15°C developed symptoms of chilling injury. In a separate study (Aina, 1990), ripening fruits were found to increase in soluble solids and carotenoid content, decrease in acidity and to undergo starch hydrolysis.

The most important product from these species (especially *I. wombolu*) is, however, the kernel of the nut, which is extracted, dried and can be stored for long periods (Plate 3). These kernels are traded on both a local and a regional scale in West Africa (£1-£3 kg<sup>-1</sup> depending on season). Uzo (1980) considered that the fruits from a single tree could generate income of US\$300 per annum. The composition of *I. gabonensis* var. *excelsa* (now *I. wombolu*) kernels at 88.1% dry matter has been reported by Ejiofor *et al.* (1987)

to be 51.3% fat, 26.0% total carbohydrate, 2.5% ash, 7.4% crude protein, 0.9% crude fibre, 9.2 mg/100g vitamin C and 0.6 mg/100g vitamin A

Other reports (eg Oke & Umoh, 1978) have quoted values of 54-67%, and even 72%, for fat content and 38.8% for carbohydrate (Ejiofor, in press). Okolo (in press) reports that the fat has an absence of volatile oils, a melting point at 37-42°C, saponification value of 233-250 and an iodine value of 2-9. He also quotes reports from 1929-39, that the myristic acid and lauric acid content of *Irvingia* kernels vary depending on the source of the fruits (Nigeria: 50.6% and 38.8%; Sierra Leone: 33.5% and 58.6% respectively). Unpublished data (Hellyer, 1997) has given myristic acid and lauric acid values of 39.2% and 51.1% from *I. wombolu* kernels from Cameroon. The amino acid composition of kernels has been reported by Amubode & Fetuga (1984).

A comparison of kernel composition between *I. gabonensis* and *I. wombolu* has shown that *I. wombolu* has less fat, more crude protein, less crude fibre and less vitamin C than *I. gabonensis* (Ejiofor *et al.*, 1987). The fat from *I. wombolu* has lower iodine and saponification values (Joseph, 1995).

Kernels are processed by grinding and separating the residue from the fat. The residue is used as a food additive to thicken soups and stews, as it produces a viscous consistency when added a few minutes before serving. A rheological study of the polysaccharides in dika nut found that the variation of 'zero-shear' specific viscosity was broadly similar to the general form of disordered polysaccharides, although with some specific attributes consistent with it having a compact molecular geometry rather than a 'random coil' conformation (Ndjouenkeu *et al.*, 1996). Joseph (1995) reports that the viscosity of mucilaginous solutions is lower at high temperatures and at high shear rates, making it appropriate as a thickening agent. The residue can be made into cubes/pellets with enhanced storage life (Ejiofor *et al.*, 1987). Okolo (in press) has calculated that a pilot plant, with a capacity of 100kg per hour, would require 256 tonnes of kernels per year. Calculated on crude protein basis, Dika nut meal shows comparatively better water and fat absorption properties than raw soy meal and hence it may have useful applications in processed foods, such as bakery products and minced meat formations (Giami, *et al.*, 1994).

## 2. *Dacryodes edulis* (G. Don) H.J. Lam and related species (African Plum, African Pear or Safoutier).

A recent workshop in Cameroon (Kengue & Nya-Ngatchou, 1994) reviewed knowledge of this species in view of new initiatives for its domestication. The flesh of the fruit has good nutritional value and has been reported by Umoro Umoti & Okiy (1987) to contain, as a percentage of dry matter (dm), 31.9% oil, 25.9% protein, 17.9% fibre. The main fatty acids in the lipid fraction are palmitic acid (36.5%), oleic acid (33.9%) and linoleic acid (24%); giving a profile similar to palm oil (*Elaeis guineensis*). The main essential amino acids are leucine (9.57%) and lysine (6.3%), while others are glutamic (17.0%), aspartic (15.1%), alanine (7.7%) acids. The ascorbic acid content of the flesh is 24.5%, but this is

lost by some forms of cooking (Achinewhu, 1983). Many of the nutrients are however in the skin of the fruit, which is usually discarded.

The seeds of *D. edulis* are usually discarded, but analysis shows them to have considerable nutritional value and a lack of toxins that makes them at least useful as a supplement to animal feed (Obasi & Okolie, 1993).

Okafor (1983) defined two varieties (*D. edulis* var. *edulis* and *D. edulis* var. *parvicarpa*) in Nigeria on the basis of their size and the relationship between their longitudinal and mid transverse circumferences. In the Congo, on the other hand, Silou (1996) characterised four fruit types that vary in size and shape. In this regard, a small survey of 3 markets in Yaoundé (Leakey & Ladipo, 1996) determined that there was 4- to 5-fold variation between fruitlots in fruit weight, pulp:seed ratio and price per kilogram of pulp (Plate 4). While large fruits with a high pulp:seed ratio were usually highly priced it was clear that some small fruits also commanded a high price, presumably because flavour, quality and other variables were important in the market place. Although Leakey & Ladipo (1996) report continuous variation in fruit size, pulp:seed ratio and other fruit characteristics, between different fruitlots of different origins, Youmbi *et al.* (1989) indicate that there are two morphological types on markets in Cameroon: a large fruit with a large seed and a small (short) fruit with a well developed mesocarp. They further indicate that these two types vary in their chemical composition, with the large type characterized by a higher lipid content in the mesocarp than in the seed, and the converse in small fruits. Fatty acid content was however not significantly different in two contrasting fruit types (Kapseu and Tchiegang, 1996). Non-structural carbohydrates are higher in the seed than in the mesocarp of both types. There is also much variation in taste and some variation in protein content (Kapseu and Tchiegang, 1996). The further characterization of these differences is important in the domestication of the species and their orientation to different markets.

Tests have determined that storage life of fruits can be prolonged beyond 8 days by refrigeration (Emebiri & Nwufu, 1990). At 15°C, storage life was two weeks, although some fruit types did deteriorate over this period. The causes of this variation in storage life need to be determined. A palm oil dip, or enclosure in a polythene bag, enhanced storage life at 15°C. At 5 °C, susceptible fruit types remained firm, but they deteriorated before day 25. The apparent genetic variation in shelflife is a trait that should be included in the selection of cultivars.

### **3. *Riciodendron heudelottii* (Baill.) Heckel (Peanut Tree, Essessang or Nyangsang).**

The kernels of the nut are widely traded in Cameroon and used as a flavouring in food dishes with the oil used in cooking (Plate 5). The paste of ground kernels is said to have a better taste than groundnut sauce (Ndoye, 1995). However, remarkably little is known about the products of this species. In Ivory Coast the kernels are used as a condiment (Téhé, 1986). The nutritive value of kernels is recorded in Pélé & Berre (1967), but the data has not been seen by the author. Kapseu (in press), however, reports that the

polyunsaturated acids are high (79.4%) and that the unsaponifiable matter is low (1.6%). The kernels can be stored for long periods.

#### 4. *Chrysophyllum albidum* G.Don (White or African Star-apple).

Achinewhu (1983) has reported that fruit pulp (Plate 6) contains 21.8mg/100g ascorbic acid, while the skin contains 75mg/100g, while Edem *et al* (1984) report 446.1 and 239.1mg/100g for pulp and skin respectively. The latter authors also indicate that proximate analysis of fruit pulp was protein (8.8%), lipid (15.1%), ash (3.4%), carbohydrate (68.7%) and crude fibre (4.0%), with only minor differences between pulp and skin. With the exception of calcium (100 v 250 mg/100g) and iron (10 v 200 mg/100g) in pulp and skin respectively, the mineral content of these components of the fruit were also very similar. According to Achinewhu (1983), the levels of toxic substances in both the mesocarp and the pericarp were low, although the juice was highly acidic. Edem *et al* (1984) on the other hand, identified high levels of tannins in pulp (627 mg/100g) and lower levels in peel (264 mg/100g). Fruit storage was best at 10°C, while for the kernel the traditional method of storing in layers of red clay was best.

The juice of fruits has potential as an ingredient of soft drinks and can be fermented for wine or other alcohol production (Ajewole & Adeyeye, 1991).

The seeds of this species are not particularly rich in lipids (3.2%), but linoleic (38.4%) and oleic (29.6%) acids are the main fatty acids present (Essien *et al.*, 1995). Ajewole & Adeyeye (1991) have, however, reported higher lipid content (16.6%) and confirmed that unsaturated fatty acids are the main component of the oil (74%) and hence desirable in the context of heart disease risk reduction. The residual cake also has potential for animal feed.

#### 5. *Garcinia kola* Heckel and related species (Bitter Cola).

The flesh of the fruit is edible and has medicinal uses (Plate 7). Comparison of the nutritive value of the pericarp and mesocarp of fresh fruits from Nigeria (Dosunmu & Johnson, 1995) shows that crude protein was higher in the mesocarp than in the pericarp (7.8% v 3.9%), while the pericarp was richer in crude fibre (16.5% v 13.9%) and macro elements (eg K: 990 v 499; Fe: 150 v 4.2; Ca 200 v 100 mg/100g). The mesocarp was richer in N (1248 v 624 mg/100g) and P (720 v 520 mg/100g). The mesocarp was also richer in crude lipid (8.7 v 6.9%) and ascorbic acid (127 v 93 mg/100g).

The kernels of the nuts are widely traded and eaten as a stimulant (Plate 8). Unsaturated fatty acids (linoleic acid: 40.5%, oleic acid: 30.8%) are the main components of the lipids (4.5%) found in the seeds of this species (Essien *et al*, 1995; Omode *et al*, 1995). The low kernel oil content of this species, however, probably eliminates it as a commercial source of oil (Foma & Abdala, 1985).

The chemical, brewing and anti-microbial properties of *Garcinia kola* seeds have been compared with hops in lager beer brewing, because of their similarity in flavour and greater availability in West Africa (Aniche & Uwakwe, 1990). Treatment of *G. kola* with methanolic lead acetate produced a yellow precipitate from which organic acids (alpha

acids) were confirmed by thin-layer chromatography. Hops, however had a higher concentration of organic acids than *G. kola*. Laboratory brewing trials with both products gave beers with similar chemical properties. Organoleptically, *G. kola* beer was as acceptable to tasters as hopped beer, but with an improved bitterness. *G. kola* and hop extracts exerted similar anti-microbial effects on two beer spoilage micro-organisms (*Candida vini* and *Lactobacillus delbruckii*).

The products of three *Garcinia* species (*G. kola* = 36%), are widely used in Ghana and 70% of this use is as chewing sticks. These are bought in urban markets as an alternative to toothpaste and brush (Adu-Tutu *et al.*, 1979). The good dental health is attributed to these chewing sticks, despite the shortage of dentists (1 per 150000 people) by comparison with the UK (1 per 3000 people), although it has to be remembered that there are also dietary differences between these countries.

## SEMI-ARID LOWLANDS OF WEST AFRICA

### 1. *Adansonia digitata* Linn. (Baobab).

The young tender leaves of Baobab are used as green or dried vegetables (Plate 9), rich in vitamin A and calcium; while the white powdery pulp of the fruit capsule is extracted and used as a flavouring in a variety of cool and hot drinks. The fruit are rich in pectins and have a vitamin C content of 169 mg/100g (Agbessi Dos-Santos 1987), at least ten-fold greater than that of oranges (Booth & Wickens, 1988). The seed kernels contain 12-15% edible oils, more protein than groundnuts and are rich in lysine, thiamine, calcium and iron (Booth & Wickens, 1988).

In the Sahel four types of Baobab are recognized. They are Black-bark, Red-bark, Grey-bark and Dark-leaf. The Dark-leaf Baobab is preferred for use as a leafy vegetable, while the Black- and Red-bark Baobabs are preferred for their fruits (Plate 10). Baobab leaf is an excellent source of calcium, iron, potassium, magnesium, manganese, molybdenum, phosphorus, and zinc (Yazzie *et al.*, 1994), and has an amino acid composition that compares favourably to that of an "ideal" protein. Dried leaves are rich in  $\beta$  carotene (Nordeide *et al.*, 1996). The leaves also contain an important amount of mucilage (Gaiwe *et al.*, 1989).

Recently, Sidibé *et al.* (1996) assessed the tree-to-tree variation in vitamin C content of fruits from the Black- Red- and Grey-bark types in 2-3 trees from 4-5 villages in three areas of Mali spanning a range of rainfall zones (450-500 mm; 600-700 mm; 750-850 mm). The vitamin C content varied 3-fold between trees, but there were no consistent differences in vitamin C content between zones or tree types. The powders from fruits are added to drinks and to gruel as it cools after cooking, so preserving the vitamins. Healthy, non-smoking adults need 23g per day of Baobab powder to meet their vitamin C requirements, while convalescents or nursing mothers require 90g.

In Senegal, Becker (1983) reported that fruit with 91.3% dry matter contained 73.7% carbohydrates, 8.9% fibre, 0.2% fat, 2.7% crude protein and 209 mg per 100g vitamin C.

In Malawi, fruit pulp of Baobab at 86.8% dry matter was found to contain: 79.4% total carbohydrate, 8.3% fibre, 4.3 % fat and 3.1% crude protein, and high levels of several minerals, including 28.4 mg g<sup>-1</sup> K and 1156µg g<sup>-1</sup> Ca (Saka *et al.*, 1994). Ascorbic acid content is 179.1 mg per 100g fresh weight (fw) (Saka, 1995). Proximate analysis of the seed kernel at 92.12% dry matter indicates that it is 29.6% fat, 28.7% crude protein and 7.3% crude fibre, while K and Ca are 1186 and 456 mg per 100g respectively. Fatty acid composition is 31.7% oleic acid, 30.8% palmitic acid and 25.2% linoleic acid.

In Nigeria, similar results were obtained (Odetokun, 1996), although carbohydrate contents were lower and protein content higher than in Malawi. Carbohydrates were higher in the pulp than in the seed and *vice versa* for protein. Eromosele *et al* (1991) observed that fruits were rich in Mg (209mg/100g) and ascorbic acid (337mg/100g).

Traditionally, Boabab seeds and pulp are sundried, roasted or fermented to extend shelflife and enhance nutritive value. When fruits from Maiduguri (Nigeria) were treated experimentally (Obizoba & Amaechi, 1993), it was found that fermentation for six days was better than roasting with regard to the value of crude protein (36.4 v 32.7%), fat (34.1 v 32.0%) and carbohydrate (30.0 v 23.5%).

Fruit pulp and aqueous extracts stored over a period of 8 months, with and without sodium metabisulphite, was found to deteriorate rapidly during periods of high humidity, unless treated with an antioxidant (Ibiyemi *et al.*, 1988). Storage of pulp was also prolonged by use of airtight containers, while juice could be stored at 10°C.

## 2. *Vitellaria paradoxa* Gaertn. syn. *Butyrospermum paradoxum* (Sheanut or Karité).

This tree is one of the most common components of the Sahelian Parklands and occurs over very large areas of Africa. The nuts (Plate 11) are used for oil extraction for cooking, soap and cosmetics. Nut production is about 3-6 kg of dry kernels per tree, but varies considerably between trees and years (Hilal, 1993; Boffa *et al*, 1996). One hundred kilograms of fruits give about 10 kg of dried kernels. These will yield about 5 kg of butter, with an oil content of 46.3-51.6% (33% non-saturated and 67% saturated).

After removing the fruit pulp, the seeds are dipped in boiling water, dried or smoked and stored. After shelling and grounding, the butter is extracted. The wet extraction process uses either boiling water or churning in cold water. The dry method uses heat and pressing. Another method uses organic solvents. The butter itself consists of a saponifiable fraction, containing triglycerides rich in vitamin F, and an unsaponifiable fraction, consisting of karitens, triterpenic alcohols, phytosterols and vitamins A, D and E, which give the butter its cosmetic hydrating, protecting, revitalizing and curative qualities (Hilal, 1993).

Many analyses have been done of Shea butter (see Booth & Wickens, 1988), but according to Sawadogo & Bezard (1982), it contains 45.6% oleic acid and 44.3% stearic acid. Oleic acid was found preferentially esterified in the 2-position (60%). The total

triacylglycerols were fractionated and the fractions were analysed for fatty acid and triglyceride compositions: the monounsaturated fraction accounted for 50% and the di-monounsaturated fraction for 27.3% of the fat. The proportion of 30 possible isomers could be determined. Only 11 isomers could be found at over 1%. Two isomers accounted for 60% of the shea butter.

According to Badifu (1989), the non-polar lipid components of shea butter were sterols, diglycerides, free fatty acids and triglycerides. The main components of non-polar lipids were triglycerides. The major fatty acids of the triglyceride were stearic acid (about 46%) and oleic acid (about 41%). Others present in relatively small quantities were 4% palmitic, 7% linoleic and 1% linolenic acids. The free sterols were 11% campesterol, 20% stigmasterol and 68% beta-sitosterol. The polar lipid components in phospholipids were phosphatidylcholine (lecithin), phosphatidylserine and phosphatidylethanolamine (cephalin). The glycolipid component was digalactosyldiglyceride and the main sugar moieties were galactose (about 32%) and glucose (about 66%). The predominant fatty acids in phospho- and glycolipids were stearic (36 to 50%), oleic (41 to 50%), and linoleic (6 to 11%).

Chavelier (1943) however reports that of the glycerides 7.0% are saturated (tributyrene 3.1%, dibutyrostearine 3.1%, arachidodipalmitine 1.0%), while 93% were non-saturated (dipalmitoleine 19%, dibutyrosoleine 54%, and palmitodioleine 19%).

After refining, traditionally prepared Shea butter is tasteless and odourless. It has been sold as baking fat, margarine and other fatty spreads and finds increasing use in edible products (Booth & Wickens, 1988). The fat is useful in patisserie and confectionary, the latex in the fat giving pliability to the dough. It is also used to formulate a cocoa butter substitute, which is unnoticeable in the final product.

Trees of *V. paradoxa* also produce a latex which can be tapped. No literature has been found giving the properties of this latex.

### **3. *Parkia biglobosa* (Jacq.) R. Br. ex G. Don (Néré or Locust Bean)**

The seeds of Néré are fermented to make Soumbala or Dawadawa, a black, strong smelling, flavoursome, tasty, proteinaceous food that is eaten for 50-90% of the year (Plate 12). This keeps without further treatment for long periods and is eaten in small quantities with sorghum or millet dumplings or porridge (Booth & Wickens, 1988). It is rich in protein (40%), lipids (35%), linoleic acid and vitamin B<sub>2</sub> (0.4-0.9 mg per 100g) and widely traded in urban markets. Soumbala is deficient in the amino acids methionine, cystine and tryptophan, like other legume seeds, but the cereals in the diet compensate for this deficiency.

The yellow, floury pulp around the seeds in the seedpod is a high energy food with up to 60% sugar (20% reducing sugars and 10-24% sucrose) and 291mg vitamin C per 100g dm (Campbell-Platt, 1980). This pulp can be eaten raw, pressed into a cake or made into a refreshing drink with water, or fermented into an alcoholic beverage. The pods and

leaves can also be eaten. Dried flour, unlike dried fermented seeds were rich in  $\forall$  and  $\exists$  carotene (Nordeide *et al.*, 1996).

#### **4. *Tamarindus indica* Linn. (Tamarind).**

Tamarind products are highly developed and widely used in Asia and so far little used in Africa, although syrup and jam are made from fruits (Plate 13). In India and Thailand especially, cultivars are grown and the food industry is active. Tamarind gum (or hydrocolloid) is a polysaccharide polymer (D-galactose, D-xylose and D-glucose) obtained from the endosperm of the seeds. It is extracted, purified and refined and used as a thickening, stabilizing and gelling agent in foods, especially in Japan where Dainippon Pharmaceutical Co conducted two years of feeding toxicity tests (Glicksman, 1986). In India it is the chief acidifying agent in curries, chutneys, and sauces. The gum can also be used as a binder in pharmaceutical tablets, as a humectant and emulifier (Hulse, 1996). Proximate analysis of seed kernels shows that 65.1-72.2% is non-fibre carbohydrate, 15.4-22.7% is protein 3.9-7.4% is oil and 0.7-8.2% is crude fibre.

Two main products are used by the food industry:

(i) Tamarind kernel powder (TKP), which contains about 50% gum and (ii) Tamarind gum polysaccharide (TGP), the purified product that is virtually 100% pure. These two products have different specifications (see Glicksman, 1986) and uses. TKP hydrates quickly in cold water, but reaches maximum viscosity if heated for 20-30 mins. TGP is more soluble but still requires some heat. A typical 1.5% gum solution will yield a viscosity of 500-800 cps at 25°C. TGP has excellent stability over a range of pH, with electrolytes (*eg.* 20% salt) and at temperatures below 65°C and degrades rapidly at higher temperatures and low pH. TSP has the ability to form gels in the presence of sugar or alcohol and can be used to form pectin-like gels in jams, jellies and other preserves (Glicksman, 1986). The xyloglucan from tamarind seeds offer no chemical advantage over guar gum as a viscosifier, but tamarind flour is cheaper indicating that a bioprocess to upgrade the tamarind polysaccharide might be commercially viable (Reid & Edwards, 1995).

In Nigeria fruits have been analysed for their ascorbic acid content (Eromosele *et al.*, 1991), but found not to be particularly rich in this vitamin.

In Malawi, tamarind fruits with 73.1% dry matter were found to contain: 85.0% total carbohydrate, 5.9% fibre, 1.6% fat, 4.1% crude protein (Saka *et al.*, 1994). Ascorbic acid content is 19.7 mg per 100g FW (Saka, 1995), but  $\forall$  and  $\exists$  carotene are absent from both dried leaves and dried fruits (Nordeide *et al.*, 1996).

#### **5. *Zizyphus mauritiana* Lam. (Jujube or Ber)**

The fruits of Jujube are one of the best edible wild fruits and some cultivars are planted. The fruits, which vary in size, are sweet and rather dry with a comparatively large stone and the larger fruits are often eaten raw (Booth & Wickens, 1988). The fruits are also

boiled with rice and millet and stewed or baked. Alternatively they are made into jellies, jams, chutneys or pickles. They can also be candied or sun-dried.

Great variation has been recorded in the fruit's nutritional value (Becker, 1983; Geurts, 1982), but they are generally rich in sugars (5.4-23%), vitamin C (96-500 mg/100g), vitamin A and carotene (21-81 mg/100g). Eromosele *et al* (1991) have reported that the fruits are rich in Ca (712.5 mg/100g) and Mg (227 mg/100g).

The potential of these fruits is virtually untapped in Africa, but is commercially exploited in India and Pakistan where cultivars are well developed – see also section below on this species under southern Africa.

## **SOUTHERN AFRICAN PLATEAU - MIOMBO WOODLANDS**

### **1. *Sclerocarya birrea* (A. Rich) Hochst. (Marula).**

The products of this pan-African, dry forest tree (fruit, nuts, oils, juice, gums, etc.) have been extensively characterized in South Africa and the findings reviewed by Weinert *et al* (1990). The fruits and nuts, in particular, have considerable commercial value. The fruits (Plate 14), which vary 30-fold in their reported flesh:stone ratio, are described as having exotic flavour and high nutritive value (*eg.* vitamin C is 2-3 times that of orange), with a few trees yielding 1.5 tonnes of fruit per tree. The strong aroma of the fruits has also characterized by freon 11 or 12 extraction and over 100 components have been identified.

The nuts too have been described as a delicacy and yield an oil with a quality (fatty acid composition) comparable with olive oil, but with a stability that is 10 times greater. This stability is explained in terms of its tocopherol /sterol composition ( $\delta^5$ -Avenasterol and  $\gamma$ -tocopherol). The amino acid content, with the exception of lysine which is deficient, has been likened to human milk and whole hens' eggs. It has been concluded that the oil could be of value to the food industry where it could be used as coating of dried fruit, as a frying oil or as a substitute for high-oleic safflower oil in baby foods.

Proximate analyses performed on fruits from different areas of southern Africa reveal some variation that may be either genetic or environmental. The causes of this variation need to be investigated as genetic variation of this magnitude would be of importance to domestication programmes. Similarly the ascorbic acid content of Marula fruits in Nigeria (403 mg/100g) has been reported to be twice that found in Botswana, although it is said to vary considerably depending on the stage of ripening, being highest in ripe fruits (Eromosele *et al.*, 1991).

The gum of Marula is acidic and has a low intrinsic viscosity, low molecular weight and high methoxyl content. The main sugar in the gum is galactose (63%), without any rhamnose (Weinert *et al.*, 1990).

White edible flesh surrounds a large nut, which contains three edible kernels. The nut represents about 50% of the weight of the fruit (Taylor & Kwerepe, 1995). In some trees the fruit pulp is sweet and in others very sour. Fruits are rich in vitamin C (194.0 mg per 100g at 85% moisture). They are very popular in Botswana and are used to make a local beer. They can vary between 10.4 and 16.0 degrees Brix (sweetness). In South Africa, the

Mitsubishi Corporation have also brewed a beer “Afreeka” which has been undergoing market trials in UK in 1997. Also in South Africa, the internationally popular liqueur, “Amarula”, is marketed by Distillers Corporation. In Zambia a wine “Marulam” is also marketed commercially (Plate 15). A pasteurised juice has also been marketed in Botswana and early ‘browning’ problems overcome (Taylor & Kwerepe, 1995). Juice flavour has been evaluated by a tasting panel, who quantified 19 characteristics of flavour, odour, mouthfeel and aftertaste. The prominent trait identified was sourness although one of five juices was much sweeter than the others (Schäfer & McGill, 1986). General experience suggests that there is considerable variation between trees within the species in sweetness, and that trees from drier environments are sweeter than those from wetter areas. Numerous small enterprises in the countries of southern Africa produce Marula jam and jellies.

The nut kernels are nutritious and widely eaten. Kernel oil is also highly prized both for cooking and for cosmetics. At 96% dry matter, the kernel is 57.3% fat, 28.3% protein, 3.7% carbohydrate and 2.9% fibre (Taylor & Kwerepe, 1995) and rich in phosphorus and magnesium. Ogbobe (1992) has however reported from Nigeria, that the kernels contain 11% crude oil, 17.2% carbohydrate, 37% crude protein, 3% fibre and 1% saponins. He also reported that the oil contains nine fatty acids of which palmitic, stearic and arachidonic acids are the most dominant.

Marula fruits fall from the tree before they are ripe, while still green, and turn yellow as they ripen on the ground. After gathering, the ripening process in storage involves concurrent changes in pH, total acids and total soluble solids. The process reaches a climax after seven days (Weinert *et al.*, 1990). Fallen fruit have a storage life of 14 days at 12.8°C. After 21 days, 89% of fruits have rotted. There are contradictory results from low temperature storage, with one report that fruits can be kept for 16 days at 4°C and another indicating that temperatures of 9°C cause damage.

Figures about the current level of production seem to be unavailable, but in 1985 it was estimated that 600 tonnes of juice were processed in South Africa. Since then there has been the introduction of several alcoholic beverages onto the international market, suggesting that the current figure must now be considerably higher. It is assumed that further growth is constrained by the fact that harvesting is restricted to the collection of fruits from wild trees, although domestication programmes have been initiated in South Africa, Botswana and Malawi.

Much research has been done on juice, extraction, product characterization and processing (see Weinert *et al.*, 1990). Variation in total soluble solids of puree and juices varied between 7.5° Brix and 15.5° Brix over three seasons, the lower value coinciding with a drought and the higher value with a wet year. Total titratable acidity was similarly affected, altering the sugar:acid ratio, an index for sensory quality in fruit juices. Sensory characteristics have also been evaluated by 14 descriptor terms; including odour, flavour, mouthfeel, aftertaste, etc. The combination of these traits and those of yield are currently being used in South Africa to register cultivars for testing horticulturally for commercial fruit production.

**2. *Uapaca kirkiana*** Muell. Arg. (Masuku or Mahobohobo).

The fleshy pulp of the Masuku fruit (Plate 16) is eaten fresh or processed into a variety of products: juices, squashes, wines, sweet beer, porridge, jams and cakes (Ngulube, 1995). In Zambia, popular brands of wine are “Masau” and “Mulunguzi”. They are produced commercially and sold in supermarkets. A beer called “Napolo Ukana” and a gin called “Kachasu” are produced.

In Zambia the fruits of Masuku are mostly (80%) cream coloured, but others are rufous (18%) and a few brown (2%) (Mwamba, 1995), with trees bearing cream coloured fruits having the greatest fruit load. The pulp forms only about 45% of the fruit, the skin being 38% and the seed 17%.

In Malawi, *Uapaca* fruits with 27.4% dry matter were found to contain: 86.5% total carbohydrate, 8.4% fibre, 1.1% fat and 1.8% crude protein (Saka *et al.* 1994). Ascorbic acid content is 16.8 mg per 100g fresh weight (Saka, 1995).

The total free sugar content of Masuku fruit juice from Zambia is 8.5% (Sufi & Kaputo, 1977) as determined by paper chromatography and confirmed by ultra violet absorption spectrophotometry. It contains glucose (4.1%), fructose (2.7%), sucrose (1.5%) and xylose (0.2%).

**3. *Zizyphus mauritiana*** Lam. (Jujube or Ber)

In Malawi, *Zizyphus* fruits at 14.8% dry matter were found to contain: 73.0% total carbohydrate, 3.4% fibre, 9.5% fat, 4.1% crude protein (Saka *et al.*, 1994). Ascorbic acid content is 13.6 mg per 100g fresh weight (Saka, 1995).

See also section above on Jujube in the Semi-arid Lowlands of West Africa.

**4. *Vangueria infausta*** Burch. (Wild Medlar).

*Vangueria* fruits in Malawi have been found to contain at 26.5% dry matter; 78.1% total carbohydrate, 10.2% fibre, 2.6% fat, 5.7% crude protein (Saka *et al.*, 1994). Ascorbic acid content is 16.8 mg per 100g fresh weight (Saka, 1995). In Botswana, ascorbic acid content of 4.7 mg per 100g has been reported for fruits with 64.4% moisture.

**5. *Azanza garckeana*** (F. Hoffm.) Exell & Hillcoat (Snotapple).

In Malawi, *Azanza* fruit pulp at 52.8% dry matter was found to contain: 35.2% total carbohydrate, 45.3% fibre, 1.1% fat, 12.0% crude protein (Saka *et al.*, 1994). Ascorbic acid content is 20.5 mg per 100g fresh weight (Saka, 1995).

## WESTERN AMAZONIA

### 1. *Inga edulis* Mart. (Inga or Guaba)

The pulp around the seeds in the pod (Plate 17) is sweet and tender and is widely marketed and eaten as a fresh fruit in Amazonia (Villachica, 1996). There is great variability within the species and potential to create cultivars, but there is not much published information on the nutritional aspects of these fruit. The pulp, which is over 80% water, is rich in carbohydrates and has high energy value. The nutritive value of fresh pulp is low, as reported by Villachica, (1996). The fruits, which can normally be kept for only 3-4 days, can be stored in a refrigerator for three weeks. The embryos of this and other *Inga* species are cooked and are more nutritious than the fruit pulp (Pennington & Robinson, in press). Boiled embryos of *Inga ilta*, for example, contain 57.7% moisture, 13.5% protein, 0.2% fat, 1.2% crude protein, 23.2% starch and 4.2% soluble carbohydrates. The cooking probably degrades trypsin inhibitors and enhances palatability.

### 2. *Bactris gasipaes* H.B.K. (Peach Palm or Pejibaye)

The two major products from Peach Palm are the fruit (mesocarp) and the 'heart of palm' (Plates 18 and 19), although the oil, wood and fibre are also valuable. The main markets for the fruit are as a delicacy for direct human consumption, as an animal feed, and as a starchy ingredient in bread and cakes. The fruit of Peach Palm, which vary in flavour and texture, is always eaten cooked as boiling breaks down a trypsin inhibitor that would otherwise have negative effects on human/animal growth. Considerable variation has, however, been reported in the presence of this inhibitor among different samples. Fruits are already marketed in jars and cans and can also be sold dehydrated.

Domestication of Peach Palm towards the different products and uses has arisen from farmer selection within Amerindian communities in tropical America (Clement 1988). Consequently, eight or nine landraces can be identified, which are suited for different uses. Classification is based on fruit size (Clement, 1990): small fruits are generally more oily and fibrous (2 landraces described); large fruits are starchy, low in oil and have a high pulp:seed ratio(2-3 landraces). The last 4 landraces are intermediate in size.

Proximate analysis of mesocarp samples have not taken into account the differences between landraces, but big variation has been reported (see review by Clement, 1990); for example, oil (8.3 - 23.0%, with one sample of 61.7%), protein (6.1-9.8%, with one sample of 17.5%), N-free extract (59.5-79.9%), fibre (2.8-9.3%). Analyses of the composition of mesocarp protein have shown that all the essential amino acids are present, although at lower levels than in maize. Arginine (7.3-9.2%) and glutamic acid (4.7-6.3%) are the most abundant. The mesocarp is frequently extremely rich in  $\beta$  carotene, although there is big variation in the presence of this pro-vitamin (Arkcoll & Aguiar, 1984).

Mesocarp oil quality has been studied in more detail than protein quality (eg. Silva & Amelotti, 1983) and contains both saturated (29.6-46.3%) and unsaturated (53.3-69.9%) fatty acids, with palmitic acid (29.6-44.8%) and oleic acid (41.0-50.3%) the most abundant respectively. It seems that the triglyceride structure is extremely variable, even within samples. This should allow opportunities for genetic selection at the clonal level. A study of the tocopherols and tocotrienols showed a strong predominance of  $\gamma$ -tocopherol (Lubrano *et al.*, 1994). Although the more primitive landraces are apparently rich in oil, there is a problem of extraction as the oil, starch and water form an emulsion that has to be solvent extracted (Clement & Arkcoll, 1985).

As an animal ration, Peach palm fruit flour can substitute for maize, sorghum or wheat and has been widely tested as a meal for chickens, usually as a partial substitute for cereals (eg 50%), especially for older birds (Clement, 1990). For pigs, silaging fruits has been reported to be an excellent means of storing the fruits, which may also be acceptable to cattle. Animal feeds would usually be based on the starchy fruit varieties with low oil content.

Peach Palm flour has also been used in bread baking and at 10% substitution for wheat gives dough with excellent baking quality (Tracy, 1996), slightly less protein, more energy (from the oil) and more vitamin A ( $\beta$ -carotene). The flour can also be used in cakes.

Palmito or Heart of Palm is already grown commercially with more than 2000ha in Costa Rica by 1990. It is, however, in competition with palmito of Acai (*Euterpe oleracea*), that has lower overheads as it is exploiting natural populations, but has lower quality control. Processing technology has been developed in Costa Rica and Brazil (Plate 20). Uses such as deep fried chips are also being found for some of the residue from palmito preparation.

## CONCLUSIONS

Farmers working with ICRAF throughout the tropics have identified the indigenous trees that they would like to see domesticated. They have also identified the traits, which from their perspective should be improved. For example they would like to see the trees coming into production at an earlier age; the length of the productive season increased; the tree height reduced and, of course, the yield and quality of the products increased. The success of this initiative to domesticate fruit trees is however closely linked to commercial (Leakey and Izac, 1996), economic and policy issues (Cannell, 1989, Leakey & Tomich, in press), but overridingly there is the need to develop and expand markets to provide the incentive to plant and manage trees in farmland. It is therefore important to examine what is known about the products and to identify ways in which they could be utilized and improved.

This review has indicated that there have been a few studies to characterize the products with commercial potential from the farmer-identified priority species. Very few of these, however, have looked at the range and origin of intraspecific genetic variation and the

opportunities it presents to improve the yield and quality of the products. Furthermore few if any of these studies have made any recommendations as to which components of the products should be improved to enhance their value to the food industry. Agroforestry researchers working in tree domestication need information from members of the food industry about the traits that they would like to see improved by genetic selection. The need is for information about characteristics that would make the products more competitive in the market, ensure their certification as a food additive, or enhance either processing or storage. Clearly dialogue and collaboration between agroforestry researchers and food scientists is needed to ensure that progress towards tree domestication is coordinated and steered in a direction that is most likely to result in the significant adoption of novel products by the food industry.

## REFERENCES

- Achinewhu, S.C. (1983). Ascorbic acid content of some Nigerian local fruits and vegetables, *Qual. Plant. Plant Foods Hum. Nutr.*, **33**, 261-266.
- Adu-Tutu, M., Afful, Y., Asante-Appiah, K., Lieberman, D., Hall, J.B. & Elvin-Lewis, M. (1979). Chewing stick usage in southern Ghana, *Econ. Bot.*, **33**, 320-328.
- Agbessi Dos-Santos, D. (1987). *Manuel de Nutrition Africaine: Elements de Base Appliquée*, Tome 1, ACCT, IPD et Editions Karthala, Dakar, Senegal.
- Aina, J.O. (1990). Physico-chemical changes in African mango (*Irvingia gabonensis*) during normal storage ripening, *Food Chem.*, **36**, 205-212.
- Ajewole, K. & Adeyeye, A. (1991). Seed oil of white star apple (*Crysophyllum albidum*) - Physicochemical characteristics and fatty acid composition, *J. Sci. Food Agric.*, **54**, 313-315.
- Akubor, P.I. (1996). The suitability of African bush mango juice for wine production, *Plant Foods Hum. Nutr.*, **49**, 213-219.
- Amubode, F.O. & Fetuga, B.L. (1984). Amino acid composition of some lesser known tree crops, *Food Chem.*, **13**, 299-307.
- Aniche, G.N. & Uwakwe, G.U. (1990). Potential use of *Garcinia kola* as substitute in lager beer brewing, *World J. Microbiol. Biotech.*, **6**, 323-327.
- Arkcoll, D.B. & Aguiar, J.P.L. (1984). Peach palm (*Bactris gasipaes* H.B.K.), a new source of vegetable oil from the wet tropics, *J. Sci. Food Agric.*, **35**, 520-526.
- Badifu, G.I.O. (1989). Lipid composition of Nigerian *Butyrospermum paradoxum* kernel, *J. Food Comp. Anal.*, **2**, 238-244.
- Becker, B. (1983). The contribution of wild plants to human nutrition in the Ferlo (Northern Senegal), *Agrofor. Syst.*, **1**, 257-267.
- Boffa, J-M., Yaméogo, G., Nikiéma, P. & Knudson, D.M. (1996). Shea nut (*Vitellaria paradoxa*) production and collection in agroforestry parklands of Burkina Faso, In: R.R.B. Leakey, A.B. Temu, M. Melnyk & P. Vantomme (eds.), *Domestication and Commercialization of Non-timber Forest Products in Agroforestry Systems, Non-Wood Forest Products No. 9*, FAO, Rome, Italy, 110-122.
- Booth F.E.M. & Wickens G.E. (1988). *Non-timber uses of selected arid zone trees and shrubs in Africa*, FAO Conservation Guide 19, FAO, Rome, Italy, 176pp.
- Cannell, M.G.R. (1989). Food crop potential of tropical trees, *Exp. Agric.*, **25**, 313-326.

- Campbell-Platt, G. (1980). African locust bean (*Parkia* species) and its West African fermented food product, dawadawa, *Ecol. Food Nutr.*, **9**, 123-132.
- Chavelier, A. (1943). Nouveau procédé de traitement des noix de Karité, *Rev. Bot. Appl.*, **5**, 536-537.
- Clement, C.R. (1988). Domestication of the pejibaye palm (*Bactris gasipaes*): past and present, *Adv. Econ. Bot.*, **6**, 155-174.
- Clement, C.R. (1990). Pejibaye, In: *Fruits of Tropical and Subtropical origin: Composition, Properties and Uses*, S. Nagy, P.E. Shaw and W.F. Wardowski (eds.), Florida Science Source Inc., Lake Alfred, Florida, USA, 302-321.
- Clement, C.R. & Arkcoll, D.B. (1985). The *Bactris gasipaes* H.B.K. (Palmae) as an oil producing crop: potential and priority of investigation, In: *Informe del Seminario-Taller Sobre Oleaginosas Promisorias*, L.E. Forero P. (ed.), Programa Interciencias de Recursos Biologicos, Bogota, Colombia.
- Dosunmu, M.I. & Johnson, E.C. (1995). Chemical evaluation of the nutritive value and changes in ascorbic acid content during storage of the fruit of 'bitter kola' (*Garcinia kola*), *Food Chem.*, **54**, 67-71.
- Ejiofor, M.A.N. (In press). Nutritional values of Ogbono (*Irvingia gabonensis* var. *excelsa*, In: D. Boland & D.O. Ladipo (eds), *Irvingia: Uses, Potential and Domestication*, ICRAF, Nairobi, Kenya, 000-000.
- Ejiofor, M.A.N., Onwubuke, S.N. & Okafor, J.C. (1987). Developing improved methods of processing and utilization of kernels of *Irvingia gabonensis* (var. *gabonensis* and var. *excelsa*), *Int. Tree Crops J.*, **4**, 283-290.
- Emebiri, L.C. & Nwufu, M.I. (1990). Effect of fruit type and storage treatments on the biodeterioration of African Pear (*Dacryodes edulis* (G. Don.) H.J. Lam.), *Int. Biodet.*, **26**, 43-50.
- Eromosele, I.C., Eromosele, C.O. & Kuzhkuzha, D.M. (1991). Evaluation of mineral elements and ascorbic acid contents in some wild plants, *Plant Foods Hum. Nutr.*, **41**, 151-154.
- Essien, E.U., Esenowo, G.J. & Akpanabiatu, M.I. (1995). Lipid composition of lesser known tropical seeds, *Plant Foods Hum. Nutr.*, **48**, 135-140.
- Foma, M. & Abdala, T. (1985). Kernel oils of seven plant species of Zaire, *J. Am. Oil Chem. Soc.*, **62**, 910-911.
- Gaiwe, R., Nkulinkiyeye-Neura, T., Bassene, E., Olschwang, D., Ba, D. & Pousset, J.L. (1989). Calcium et mucilage dans les feuilles de *Adansonia digitata* (Boabab), *Int. J. Crude Drug Res.*, **27**, 101-104.
- Geurts, I.F. (1982). The Indian jujube or ber (*Zizyphus mauritiana* Lamk.). Aspects related to germplasm conservation. *A Preliminary Report*, Royal Tropical Institute, Amsterdam, The Netherlands.
- Giami, S.Y., Okonkwo, V.I. & Akusu, M.O. (1994). Chemical composition and functional properties of raw, heat-treated and partially proteolysed wild mango (*Irvingia gabonensis*) seed flour, *Food Chem.*, **49**, 237-243.
- Glicksman, M. (1996). Tamarind seed gum, In: M. Glicksman (Ed.), *Food Hydrocolloids, Volume III*, CRC Press Inc., Boca Raton, Florida, USA, 191-202.
- Hilal, N.S. (1993). Contribution à l'étude du Karité, *Butyrospermum paradoxum* (Gaertn.F.) Hepper Sapotaceae), Monograph of Faculté de Medecine et de Pharmacie, University of Cheikh Anta Diop, Dakar, Senegal, 144p.

- Hulse, J. (1996). Flavours, spices and edible gums: opportunities for integrated agroforestry systems, R.R.B. Leakey, A.B. Temu, M.Melnyk & P. Vantomme (eds.), *Domestication and Commercialization of Non-timber Forest Products in Agroforestry Systems, Non-Wood Forest Products No. 9*, FAO, Rome, Italy, 86-96.
- Ibiyemi, S.A., Abiodun, A. & Akanji, S.A. (1988). *Andersonia [sic] digitata* Bombax and *Parkia filicoideae* Welw: Fruit pulp for the soft drink industry, *Food Chem.*, **28**, 111-116.
- ICRAF (1997). ICRAF Medium-Term Plan 1998-2000, ICRAF, PO Box 30677, Nairobi, Kenya, 73p.
- Joseph, J.K. (1995). Physico-chemical attributes of wild mango (*Irvingia gabonensis*) seeds, *Biores. Tech.*, **53**, 179-181.
- Joseph, J.K. & Aworh, O.C. (1991). Composition, sensory quality and respiration during ripening and storage of edible wild mango (*Irvingia gabonensis*), *Int. J. Food Sci. Tech.*, **26**, 337-342.
- Kapseu, C. (In press). Composition en acides gras et en triglycérides des huiles des oléagineux non conventionnels, In: *Deuxième Séminaire International sur la Valorisation du Safoutier et autres Oléagineux non Conventionnels*, C Kapseu and G.J. Kayem (eds), Ngaoundere, Cameroon, 3-5 December, 1997.
- Kapseu, C and Tchiegang, C. (1996). Composition de l'huile des fruits de deux cultivars de safou au Cameroun, *Fruits*, **51**, 185-191.
- Kengue, J. & Nya-Ngatchou, J. (1994). Le Safoutier: the African pear. *Proceedings of the Domestication of the African Pear Workshop*, Douala, Cameroon, 4-6 October 1994.
- Ladipo, D.O. Fondoun, J-M. & Ganga, N. (1996). Domestication of the bush mango (*Irvingia* spp.): some exploitable intraspecific variations in west and central Africa, In: R.R.B. Leakey, A.B. Temu, M.Melnyk & P. Vantomme (eds.), *Domestication and Commercialization of Non-timber Forest Products in Agroforestry Systems, Non-Wood Forest Products No. 9*, FAO, Rome, Italy,, 193-205.
- Leakey, R.R.B. & Izac, A-M.N. (1996). Linkages between domestication and commercialization of non-timber forest products: implications for agroforestry, In: R.R.B. Leakey, A.B. Temu, M.Melnyk & P. Vantomme (Eds.), *Domestication and Commercialization of Non-timber Forest Products in Agroforestry Systems, Non-Wood Forest Products No. 9*, FAO, Rome, Italy, 1-7.
- Leakey, R.R.B. & Ladipo, D.O. (1996). Trading on genetic variation - fruits of *Dacryodes edulis*, *Agrofor. Today*, **8**(2), 16-17.
- Leakey, R.R.B. & Simons, A.J. (1997). The domestication and commercialization of indigenous trees in agroforestry for the alleviation of poverty, *Agrofor. Sys.*, **0**, 000-000.
- Leakey, R.R.B. & Tomich, T.P. (In press). Domestication of tropical trees: from biology to economics and policy, In: L.E. Buck, J.P. Lassoie & E.C.M. Fernandes (eds.), *Agroforestry in Sustainable Ecosystems*, CRC Press/Lewis Publishers, New York, USA.
- Lubrano, C., Robin, J.R. & Khaiat, A. (1994). Composition en acides gras, stérols et tocophérols d'huiles de pulpe de fruits de six espèces de palmiers de Guyane, *Oleagineux*, **49**, 59-65.
- Mwamba, C.K. (1995). Natural variation in fruits of *Uapaca kirkiana* in Zambia, *For. Ecol. Manage.*, **26**, 299-303.

- Ndjouenkeu R., Goycoolea, F.M., Morris, E.R. & Akingbala, J.O. (1996). Rheology of okra (*Hibiscus esculentus* L.) and dika nut (*Irvingia gabonensis*) polysaccharides, *Carbohydr. Polym.*, **29**, 263-269.
- Ndoye, O. (1995). The markets for non-timber forest products in the humid forest zone of Cameroon and its borders: Structure, conduct, performance and policy implications, *Report to CIFOR*, Bogor, Indonesia, 86 p.
- Ngulube, M. R. (1995). Indigenous fruit trees in southern Africa: the potential of *Uapaca kirkiana*, *Agrofor. Today*, **7**(3-4), 17-18.
- Nordeide, M.B., Hatlpy, A., Folling, M. Lied, E. & Oshaug, A. (1996). Nutrient composition and nutritional importance of green leaves and wild food resources in an agricultural district, Koutiala, in Southern Mali, *Int. J. Food Sci. Nutr.*, **47**, 455-468.
- Obasi, N.B.B. & Okolie, N.P. (1993). Nutritional constituents of the seeds of the African pear, *Dacryodes edulis*, *Food Chem.*, **46**, 297-299.
- Obizoba, I.C. & Amaechi, N.A. (1993). The effect of processing methods on the chemical composition of baobab (*Adansonia digitata*.L.) pulp and seed, *Ecol. Food Nutr.*, **29**, 199-205.
- Odetokun, S.M. (1996). The nutritive value of baobab fruit (*Adansonia [sic] digitata*), *Riv. Ital. Sost. Gras.*, **23**, 371-373.
- Ogbobe, O. (1992). Physicochemical composition and characterization of the seed and oil of *Sclerocarya birrea*, *Plant Foods Hum. Nutr.*, **42**, 201-206.
- Okafor, J.C. (1983). Varietal delimitation in *Dacryodes edulis* (G.DON) H.J. LAM (Burseraceae), *Int. Tree Crops J.*, **2**, 255-265.
- Oke, O.L. & Umoh, I.B. (1978). Lesser known oilseeds.I. Chemical composition, *Nutr. Rep. Int.*, **17**, 293-297.
- Okolo, H.C. (In press). Industrial potential of various *Irvingia gabonensis* products such as oil, Ogbono and juice, In: D. Boland & D.O. Ladipo (eds), *Irvingia: Uses, Potential and Domestication*, ICRAF, Nairobi, Kenya, 000-000.
- Omode, A.A., Fatoki, O.S. & Olaogun, K.A. (1995). Physicochemical properties of some underexploited and nonconventional oilseeds, *J. Agric. Food Chem.*, **43**, 2850-2853.
- Pélé J. & Berre, S. (1967). Les aliments d'origine végétale au Camérout, *Cam. Agric., Past. For.*, **108-111**, ORSTOM, Paris, France.
- Pennington, T.D. & Robinson, R.K. (In press). Utilization profile of a new species: *Inga ilta* T.D. Penn., In: T.D. Pennington and E. Fernandes (Eds.), *The Genus Inga: Utilization*, Royal Botanic Garden, Kew, UK and ICRAF, Nairobi, Kenya.
- Reid, J.S.G. & Edwards, M.E. (1995). In: A.M. Stephen, 155-186, *Food Polysaccharides and their Applications*, Marcel Dekker Inc., New York, USA.
- Saka J.D.K. (1995). The nutritional value of edible indigenous fruits: present research status and future directions, In: J.A. Maghembe, Y. Ntupanyama & P.W.Chirwa (eds.), *Improvement of Indigenous Fruit Trees of the Miombo Woodlands of Southern Africa*, ICRAF, PO Box 30677, Nairobi, Kenya, 50-57.
- Saka, J.D.K., Msonthi, J.D. & Maghembe, J.A. (1994). Nutritional value of edible fruits of indigenous wild trees in Malawi, *Forest Ecology and Management*, **64**, 245-248.
- Sawadogo, K. & Bezar, J. (1982). Etude de la structure glyceridique du beurre de karité, *Oleagineux*, **37**, 69-74.

- Schäfer, G. & McGill, A.E.J. (1986). Flavour profiling of juice of the Marula (*Sclerocarya birrea* subsp. *caffra*) as an index for cultivar selection, *Acta Hort.*, **194**, 215-222.
- Sidibé, M., Scheuring, J.F., Tembely, D., Sidibé, M.M., Hofman, P. & Frigg, M. (1996). Boabab – homegrown vitamin C for Africa, *Agrofor. Today*, **8** (2), 13-15.
- Silou, T. (1996). Le safoutier (*Dacryodes edulis*): un arbre mal connu, *Fruits*, **51**, 47-60.
- Silva W.G. & Amelotti G. (1983). Composition of the fatty substances from the fruit of *Guiljelma speciosa* (Pupunha). *Riv. Ital. Sost. Gras.*, **60**, 767-770.
- Sufi, N.A. & Kaputo, M.T. (1977). Identification and determination of free sugars in Masuku fruit (*Uapaca kirkiana*), *Zam. J. Sci. Tech.*, **2**, 23-25.
- Taylor, F.W. & Kwerepe, B. (1995). Towards domestication of some indigenous fruit trees in Botswana, In: J.A. Maghembe, Y. Ntupanyama and P.W.Chirwa (eds.), *Improvement of Indigenous Fruit Trees of the Miombo Woodlands of Southern Africa*, ICRAF, PO Box 30677, Nairobi, Kenya, 113-134.
- Téhé, H. (1986). Utilizations des ressources forestières chez les Guérés et les Oubis Côte d'Ivoire), *Banco, (Côte d'Ivoire)*, **4**, 26-30.
- Tracy, M.D. (1996). Pejibaye flour, a hopeful option, *Bol. Retadar*, **23**, 3.
- Umoro Umati, U. & Okiy, A. (1987). Characteristics and composition of the pulp oil and cake of the African pear, *Dacryodes edulis* (G.Don) H.J.Lam., *J. Sci.Food Agric.*, **38**, 67-72.
- Uzo, J.O. (1980). Yield and harvest predictions of some indigenous perennial fruits, roots and leafy vegetables in tropical West Africa, *Proceedings of International Symposium on the Current Problems of Fruits and Vegetables (Tropical and Sub-tropical)*, Laguna, 24-26 March 1980, 22p.
- Villachica, H. (1996). *Frutales y Hortalizas Promisorios de la Amazonia*, Tratado de Cooperacion Amazonica, Lima, Peru, 367p.
- Weinert, I.A.G., van Wyk, P.J. & Holtzhausen, L.C. (1990). Marula, In: *Fruits of Tropical and Subtropical origin: Composition, Properties and Uses*, S. Nagy, P.E. Shaw and W.F. Wardowski (eds.), Florida Science Source Inc., Lake Alfred, Florida, USA, 88-115.
- Yazzie, D., VanderJagt, D.J., Pastuszyn, A., Okolo, A. & Glew, R.H. (1994). The amino acid and mineral content of baobab (*Adansonia digitata* L.) leaves, *J. Food Comp. Anal.*, **7**, 189-193.
- Youmbi, E., Clair-Maczulajtys & Bory, G. (1989). Variations de la composition chimique des fruits de *Dacryodes edulis* (DON) LAM., *Fruits*, **44**, 149-153.